

WHAT IS CLAIMED IS:

1. A process for detecting a target single-stranded nucleic acid having a first base sequence, said process comprising the steps of:

forming a double-stranded nucleic acid by hybridizing said target single-stranded nucleic acid with a probe nucleic acid having a second base sequence complementary to said first base sequence;

providing a chemiluminescent compound capable of being associated with a double-stranded nucleic acid, and then associating said chemiluminescent compound with the double-stranded nucleic acid resulting from said forming step; and

detecting luminescence from said chemiluminescent compound associated with said double-stranded nucleic acid.

2. The process according to Claim 1, wherein the luminescence-detecting step is conducted under a condition that only said chemiluminescent compound associated with said double-stranded nucleic acid can exhibit chemiluminescence.

3. The process according to Claim 2, wherein said condition is in an aqueous medium in which said chemiluminescent compound non-associated with a double-

TE 2700 "040704"

stranded nucleic acid does not exhibit chemiluminescence.

4. The process according to Claim 3, wherein said aqueous medium is water.

5. The process according to Claim 3, wherein said aqueous medium is an aqueous buffer solution.

6. The process according to Claim 3, wherein said aqueous medium is a mixture solution of water and an organic solvent miscible with water.

7. The process according to Claim 6, wherein said organic solvent comprises at least one solvent selected from the group consisting of methanol, ethanol, acetonitrile, dimethylformamide, dimethylsulfoxide, and isopropanol.

8. The process according to Claim 6, wherein said mixture solution has a content of said organic solvent falling within 2 to 50% by volume relative to water.


9. The process according to Claim 8, wherein said content falls within 5 to 20% by volume relative to water.

10. The process according to Claim 3, wherein pH of

said aqueous medium ranges from 5 to 8.

11. The process according to Claim 1, wherein said chemiluminescent compound is capable of being inserted into the double helical structure of said double-stranded nucleic acid as an intercalator.

12. The process according to Claim 11, wherein said chemiluminescent compound is a pyrylium compound represented by the following formula [1]:



[1],

wherein:

X is O, S, Se or Te;

two of R¹, R² and R³ are independently a substituted or unsubstituted aryl group;



the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:

L is -L¹-, -L²-L³- or -L⁴-L⁵-L⁶-, wherein each of

102730" 2401E60


A is a substituted or unsubstituted aryl group, or $-CH=R^5$, wherein R^5 is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and

Y^- is an anion.

$$\begin{array}{c} -\text{CH}=\text{C}- (\text{CH}=\text{CH})_n- \\ | \\ \text{Z} \end{array} \quad [2];$$
$$-\text{CH}=\text{CH}-\phi-\text{CH}=\text{CH}- \quad [4];$$


wherein Z is a hydrogen atom or a substituted or unsubstituted lower alkyl group, n is 0, 1 or 2, and ϕ is a substituted or unsubstituted o-, m- or p-phenylene group.


14. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [7]:



wherein X is O, S, Se, or Te, and Y is an anion.

15. The process according to Claim 14, wherein X is O or S, and Y is I or ClO_4 .

16. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [8]:




[8],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

17. The process according to Claim 16, wherein X is O or S, and Y is I or ClO₄.

18. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [9]:




wherein X is O, S, Se, or Te, and Y⁻ is an anion.

19. The process according to Claim 18, wherein X is O

or S, and Y is I or ClO₄.

20. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [10]:




[10],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

21. The process according to Claim 20, wherein X is O or S, and Y is I or ClO₄.


22. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [11]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

23. The process according to Claim 22, wherein X is O or S, and Y is I or ClO₄.


24. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [12]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

25. The process according to Claim 24, wherein X is O or S, and Y is I or ClO₄.

26. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [13]:




[13],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

27. The process according to Claim 26, wherein X is O or S, and Y is I or ClO₄.


28. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [14]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

29. The process according to Claim 28, wherein X is O or S, and Y is I or ClO₄.

30. The process according to Claim 12, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [15]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

T02F00" 8107E660

31. The process according to Claim 30, wherein X is O or S, and Y is I or ClO₄.

32. The process according to Claim 12, wherein at least one hydrophilic group is introduced into at least one substituent of said pyrylium compound.

33. The process according to Claim 1, wherein said chemiluminescent compound is inserted into said double-stranded nucleic acid by groove binding.

34. The process according to Claim 1, wherein said luminescence-detecting step includes allowing said chemiluminescent compound and said double-stranded nucleic acid to coexist with an oxalic ester and hydrogen peroxide.

35. The process according to Claim 34, wherein said oxalic ester is bisdinitrophenyl oxalate.

36. The process according to Claim 1, wherein said step of forming a double-stranded nucleic acid includes immobilizing said target nucleic acid or said probe nucleic acid to a solid phase previous to hybridizing said target nucleic acid with said probe nucleic acid.

00931016-091704

37. The process according to Claim 1, wherein said step of forming a double-stranded nucleic acid includes the steps of:

immobilizing said target nucleic acid to a solid phase;
preparing as said probe nucleic acid, a single-stranded nucleic acid capable of binding with said target nucleic acid through complementary sequences at 3'-end regions of the target nucleic acid and the single-stranded nucleic acid;

hybridizing said target nucleic acid with said probe nucleic acid to form a double-stranded nucleic acid; and

polymerizing nucleotides to the 3'-ends of said target nucleic acid and said probe nucleic acid by extension reaction to extend a double-stranded portion in said double-stranded nucleic acid.

38. The process according to Claim 1, wherein said step of forming a double-stranded nucleic acid includes the steps of:

preparing as said probe nucleic acid, a single-stranded nucleic acid capable of binding with said target nucleic acid through complementary sequences at 3'-end regions of the target nucleic acid and the single-stranded nucleic acid;

39. The process according to Claim 36, 37 or 38, wherein said solid phase comprises a plastic plate.

41. The process according to claim 40, wherein said DNA is cDNA.

43. The process according to Claim 1, wherein said probe nucleic acid is DNA or RNA.

44. The process according to Claim 1, wherein said target nucleic acid is mRNA comprising a base sequence

adenylic ac
nucleic acid
oligodeoxyr
rid, the bas
rget nuclei
uantifying a
t base sequ
unded nuclei
d nucleic a
nd base seq
nescent com
tranded nuc
nescent com
lting from
e from said
aid double-
rding to Cl
p is conduc
scent compo
ic acid can

measuring luminescence from said chemiluminescent compound associated with said double-stranded nucleic acid.

46. The process according to Claim 45, wherein the luminescence-measuring step is conducted under a condition that only said chemiluminescent compound associated with said double-stranded nucleic acid can exhibit chemiluminescence.

48. The process according to Claim 47, wherein said aqueous medium is water.

50. The process according to Claim 47, wherein said aqueous medium is a mixture solution of water and an organic solvent miscible with water.

52. The process according to Claim 50, wherein said mixture solution has a content of said organic solvent falling within 2 to 50% by volume relative to water.

[illegible][illegible]

According to [1]:

d is capable of forming a pyridine-like ring.

The structure of said compound is shown below:

[1]:

R²

X⁺

Y⁻


R³

are independent of each other;


where R¹ is a hydrogen atom,

R² is a methyl group,

According to the present invention, the compound is a pyridine derivative of the following structure [1]:



where R² and R³ are independently hydrogen, alkyl, alkenyl, aryl, or heteroaryl groups; X is a halogen atom, a cyano group, or a nitro group; and Y⁻ is a halogen atom, a cyano group, or a nitro group.



are independent
;
oxygen atom,
methyl group,

re independent
;
oxygen atom,
yl group,

re independent
;
oxygen atom,
methyl group,

re independent
;
hydrogen atom,
methyl group,

oxygen atom,
methyl group,



group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:


L is $-L^1-$, $-L^2-L^3-$ or $-L^4-L^5-L^6-$, wherein each of L^1 to L^6 is independently $-(CH=CH)-$, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene group, or $-CH=R^4-$, wherein R^4 is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or $-CH=R^5$, wherein R^5 is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and


Y^- is an anion.

57. The process according to Claim 56, wherein L in said formula [1] is any one of the groups represented by the following formulae [2] to [6], respectively:






[5]; and



[6],

wherein Z is a hydrogen atom or a substituted or unsubstituted lower alkyl group, n is 0, 1 or 2, and ϕ is a substituted or unsubstituted o-, m- or p-phenylene group.

58. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [7]:




[7],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

59. The process according to Claim 58, wherein X is O or S, and Y is I or ClO₄.


60. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [8]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

61. The process according to Claim 60, wherein X is O or S, and Y is I or ClO₄.


62. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [9]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

63. The process according to Claim 62, wherein X is O or S, and Y is I or ClO₄.

64. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [10]:




wherein X is O, S, Se, or Te, and Y⁻ is an anion.

65. The process according to Claim 64, wherein X is O or S, and Y is I or ClO₄.

"FOOTED" SHOWS

66. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [11]:




[11],

wherein X is O, S, Se, or Te, and Y^- is an anion.

67. The process according to Claim 66, wherein X is O or S, and Y is I or ClO_4 .


68. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [12]:



wherein X is O, S, Se, or Te, and Y is an anion.

69. The process according to Claim 68, wherein X is O or S, and Y is I or ClO₄.


70. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [13]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

71. The process according to Claim 70, wherein X is O or S, and Y is I or ClO₄.


72. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [14]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

73. The process according to Claim 72, wherein X is O or S, and Y is I or ClO₄.

74. The process according to Claim 56, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [15]:



[15],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

75. The process according to Claim 74, wherein X is O or S, and Y is I or ClO₄.

76. The process according to Claim 56, wherein at least one hydrophilic group is introduced into at least one substituent of said pyrylium compound.

77. The process according to Claim 45, wherein said chemiluminescent compound is inserted into said double-stranded nucleic acid by groove binding.

78. The process according to Claim 45, wherein said luminescence-measuring step includes allowing said chemiluminescent compound and said double-stranded nucleic acid to coexist with an oxalic ester and hydrogen peroxide.

82. The process according to Claim 45, wherein said step of forming a double-stranded nucleic acid includes the steps of:

preparing as said probe nucleic acid, a single-stranded nucleic acid capable of binding with said target nucleic acid through complementary sequences at 3'-end regions of the target nucleic acid and the single-stranded nucleic acid;

immobilizing said probe nucleic acid to a solid phase;

hybridizing said target nucleic acid with said probe nucleic acid to form a double-stranded nucleic acid; and

polymerizing nucleotides to the 3'-ends of said target nucleic acid and said probe nucleic acid by extension reaction to extend a double-stranded portion in said double-stranded nucleic acid.

83. The process according to Claim 80, 81 or 82, wherein said solid phase comprises a plastic plate.

84. The process according to Claim 45, wherein said target nucleic acid is DNA or RNA.

85. The process according to Claim 84, wherein said DNA is cDNA.

009304000000

86. The process according to Claim 84, wherein said RNA is mRNA, tRNA or rRNA.

87. The process according to Claim 45, wherein said probe nucleic acid is DNA or RNA.

88. The process according to Claim 45, wherein said target nucleic acid is mRNA comprising a base sequence corresponding to oligoriboadenylic acid at 3'-end region thereof, and said probe nucleic acid comprises a base sequence corresponding to oligodeoxyribothymidylic acid or polydeoxyribothymidylic acid, the base sequence relating to hybridization with said target nucleic acid.

89. A process for detecting a target double-stranded nucleic acid comprising the steps of:

providing a chemiluminescent compound capable of being associated with a double-stranded nucleic acid, and then associating said chemiluminescent compound with the target double-stranded nucleic acid; and

detecting luminescence from said chemiluminescent compound associated with said target double-stranded nucleic acid.

90. The process according to Claim 89, wherein the


96. The process according to Claim 94, wherein said mixture solution has a content of said organic solvent falling within 2 to 50% by volume relative to water.

97. The process according to Claim 96, wherein said content falls within 5 to 20% by volume relative to water.

98. The process according to Claim 91, wherein pH of said aqueous medium ranges from 5 to 8.

99. The process according to Claim 89, wherein said chemiluminescent compound is capable of being inserted into the double helical structure of said double-stranded nucleic acid as an intercalator.

100. The process according to Claim 99, wherein said chemiluminescent compound is a pyrylium compound represented by the following formula [1]:



[1],

wherein:

X is O, S, Se or Te;

two of R¹, R² and R³ are independently a substituted or unsubstituted aryl group;


the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:




L is -L¹-, -L²-L³- or -L⁴-L⁵-L⁶-, wherein each of L¹ to L⁶ is independently -(CH=CH)-, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene group, or -CH=R⁴-, wherein R⁴ is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or -CH=R⁵, wherein R⁵ is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and

Y⁻ is an anion.


101. The process according to Claim 100, wherein L in said formula [1] is any one of the groups represented by the following formulae [2] to [6], respectively:





wherein Z is a hydrogen atom or a substituted or unsubstituted lower alkyl group, n is 0, 1 or 2, and ϕ is a substituted or unsubstituted o-, m- or p-phenylene group.

102. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [7]:




T02F00"840TE060

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

103. The process according to Claim 102, wherein X is O or S, and Y is I or ClO₄.

104. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [8]:




[8],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

105. The process according to Claim 104, wherein X is O or S, and Y is I or ClO₄.


106. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [9]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

107. The process according to Claim 106, wherein X is O or S, and Y is I or ClO₄.


108. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [10]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

109. The process according to Claim 108, wherein X is O or S, and Y is I or ClO₄.

110. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [11]:




[11],

wherein X is O, S, Se, or Te, and Y^- is an anion.

111. The process according to Claim 110, wherein X is O or S, and Y is I or ClO_4 .

112. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [12]:


T02780" CHOT E660



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

113. The process according to Claim 112, wherein X is O or S, and Y is I or ClO₄.

114. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [13]:




[13],

"SECRET"

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

115. The process according to Claim 114, wherein X is O or S, and Y is I or ClO₄.


116. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [14]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

117. The process according to Claim 116, wherein X is O or S, and Y is I or ClO₄.

118. The process according to Claim 100, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [15]:



[15],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

119. The process according to Claim 118, wherein X is O or S, and Y is I or ClO₄.

120. The process according to Claim 100, wherein at least one hydrophilic group is introduced into at least one substituent of said pyrylium compound.

121. The process according to Claim 89, wherein said luminescence-detecting step includes allowing said chemiluminescent compound and said double-stranded nucleic acid to coexist with an oxalic ester and hydrogen peroxide.

122. The process according to Claim 121, wherein said oxalic ester is bisdinitrophenyl oxalate.

123. The process according to Claim 89, wherein said

127. The process according to Claim 126, wherein said

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume

process according to

in ranges from 5

process according to

water.

process according to

an aqueous buffer

process according to

a mixture solution

with water.

process according to

comprises at least

ing of methanol,

dimethylsulfoxide

process according to

as a content of

to 50% by volume

process according to

in 5 to 20% by volume


process according to

in ranges from 5

process according to

chemiluminescent compound is capable of being inserted into the double helical structure of said double-stranded nucleic acid as an intercalator.

135. The process according to Claim 134, wherein said chemiluminescent compound is a pyrylium compound represented by the following formula [1]:



wherein:

X is O, S, Se or Te;

two of R¹, R² and R³ are independently a substituted or unsubstituted aryl group;

the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:

L is -L¹-, -L²-L³- or -L⁴-L⁵-L⁶-, wherein each of L¹ to L⁶ is independently -(CH=CH)-, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene




T 0 2 3 0 " S 1 0 7 E 6 6 0

group, or $-\text{CH}=\text{R}^4-$, wherein R^4 is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or $-\text{CH}=\text{R}^5$, wherein R^5 is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and

Y^- is an anion.

136. The process according to Claim 135, wherein L in said formula [1] is any one of the groups represented by the following formulae [2] to [6], respectively:




wherein Z is a hydrogen atom or a substituted or unsubstituted lower alkyl group, n is 0, 1 or 2, and ϕ is a

FOI 800 807660

substituted or unsubstituted o-, m- or p-phenylene group.


137. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [7]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

138. The process according to Claim 137, wherein X is O or S, and Y is I or ClO₄.

139. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [8]:




[8],

wherein X is O, S, Se, or Te, and Y^- is an anion.

140. The process according to Claim 139, wherein X is O or S, and Y is I or ClO_4 .

141. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [9]:




[9],

wherein X is O, S, Se, or Te, and Y^- is an anion.

TOP SECRET

142. The process according to Claim 141, wherein X is O or S, and Y is I or ClO₄.

143. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is a compound represented by the following formula [10]:




[10],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

144. The process according to Claim 143, wherein X is O or S, and Y is I or ClO₄.


145. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [11]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

146. The process according to Claim 145, wherein X is O or S, and Y is I or ClO₄.


147. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [12]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

148. The process according to Claim 147, wherein X is O or S, and Y is I or ClO₄.


149. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [13]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

150. The process according to Claim 149, wherein X is O or S, and Y is I or ClO₄.


151. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [14]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

152. The process according to Claim 151, wherein X is O or S, and Y is I or ClO₄.

153. The process according to Claim 135, wherein said chemiluminescent compound represented by said formula [1] is the compound represented by the following formula [15]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

154. The process according to Claim 153, wherein X is O or S, and Y is I or ClO₄.

155. The process according to Claim 135, wherein at least one hydrophilic group is introduced into at least one substituent of said pyrylium compound.

156. The process according to Claim 124, wherein said luminescence-measuring step includes allowing said chemiluminescent compound and said double-stranded nucleic acid to coexist with an oxalic ester and hydrogen peroxide.

157. The process according to Claim 156, wherein said oxalic ester is bisdinitrophenyl oxalate.

158. The process according to Claim 124, wherein said chemiluminescent compound is inserted into said double-stranded nucleic acid by groove binding.

159. A process for detecting a target single-stranded nucleic acid having a first base sequence, said process comprising the steps of:

forming a double-stranded nucleic acid by hybridizing said target single-stranded nucleic acid with a probe

providing a compound which can be intercalated into a double-stranded nucleic acid and which is capable of exhibiting chemiluminescence only in a hydrophobic condition, and then intercalating said compound into the double-stranded nucleic acid resulting from said forming step; and

160. A process for quantifying a target single-stranded nucleic acid having a first base sequence, said process comprising the steps of:

forming a double-stranded nucleic acid by hybridizing said target single-stranded nucleic acid with a probe nucleic acid having a second base sequence complementary to said first base sequence;

providing a compound which can be intercalated into a double-stranded nucleic acid and which is capable of exhibiting chemiluminescence only in a hydrophobic condition, and then intercalating said compound into the

did resulting from
s medium said doubt
aid compound is in
apable of causing
, and measuring the

detecting a target
ne steps of:

which can be inter
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doub
aid compound is in
apable of causing
, and measuring th

detecting a target
ne steps of:

which can be inte
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doub
aid compound is in
apable of causing
, and measuring th

detecting a target
ne steps of:

which can be inte
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doub
aid compound is in
apable of causing
, and measuring th

detecting a target
ne steps of:

which can be inte
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doubt
aid compound is in
apable of causing
, and measuring the

detecting a target
ne steps of:

which can be inter
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doub
aid compound is in
apable of causing
, and measuring th

detecting a target
ne steps of:

which can be inte
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

did resulting from
s medium said doub
aid compound is in
apable of causing
, and measuring th

detecting a target
ne steps of:

which can be inte
aid and which is c
emiluminescence on
l then intercalati
randed nucleic ac
s medium said doub
aid compound is in
pable of causing s
and detecting the

uantifying a target
rising the steps c
which can be inter
aid and which is ca

placing in an aqueous medium said double-stranded nucleic acid into which said compound is intercalated together with a reagent capable of causing said compound to exhibit chemiluminescence, and measuring the resulting chemiluminescence.

following formula [1]:

[1],

X is O, S, Se or Te;

the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted


cycloalkyl group, -A or -L-A, wherein:

L is $-L^1-$, $-L^2-L^3-$ or $-L^4-L^5-L^6-$, wherein each of L^1 to L^6 is independently $-(CH=CH)-$, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene group, or $-CH=R^4-$, wherein R^4 is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or $-CH=R^5$, wherein R^5 is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and

Y^- is an anion.

164. The quantifying process according to Claim 160 or 162, wherein said compound is a pyrylium compound represented by the following formula [1]:



wherein:

X is O, S, Se or Te;

two of R^1 , R^2 and R^3 are independently a substituted or

T-02-F-03-0107-0000

unsubstituted aryl group;


the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:

L is -L¹-, -L²-L³- or -L⁴-L⁵-L⁶-, wherein each of L¹ to L⁶ is independently -(CH=CH)-, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene group, or -CH=R⁴-, wherein R⁴ is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or -CH=R⁵, wherein R⁵ is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and

Y⁻ is an anion.

165. A pyrylium compound represented by the following formula [1] for use in chemiluminescence analysis:



wherein:

X is O, S, Se or Te;

two of R¹, R² and R³ are independently a substituted or unsubstituted aryl group;

the other is a hydrogen atom, halogen atom, sulfonate group, amino group, styryl group, nitro group, hydroxyl group, carboxyl group, cyano group, substituted or unsubstituted alkyl group, substituted or unsubstituted cycloalkyl group, -A or -L-A, wherein:

L is -L¹-, -L²-L³- or -L⁴-L⁵-L⁶-, wherein each of L¹ to L⁶ is independently -(CH=CH)-, a divalent group derived from the substituted or unsubstituted aryl group, a substituted or unsubstituted lower alkylene group, or -CH=R⁴-, wherein R⁴ is a ring structure having an oxo group; and

A is a substituted or unsubstituted aryl group, or -CH=R⁵-, wherein R⁵ is a substituted or unsubstituted heterocyclic ring, substituted or unsubstituted cycloalkyl group or substituted or unsubstituted aromatic ring; and


Y⁻ is an anion.

166. The pyrylium compound according to Claim 165, wherein L in said formula [1] is any one of the groups represented by the following formulae [2] to [6],

0991048-08704
102780-8107860

wherein Z is a hydrogen atom or a substituted or unsubstituted lower alkyl group, n is 0, 1 or 2, and ϕ is a substituted or unsubstituted o-, m- or p-phenylene group.


represented by the following formula [7]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

168. The pyrylium compound according to Claim 167, wherein X is O or S, and Y is I or ClO₄.

169. The pyrylium compound according to Claim 165, represented by the following formula [8]:




[8],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

170. The pyrylium compound according to Claim 169, wherein X is O or S, and Y is I or ClO₄.

171. The pyrylium compound according to Claim 165, represented by the following formula [9]:




[9],

wherein X is O, S, Se, or Te, and Y- is an anion.

172. The pyrylium compound according to Claim 171, wherein X is O or S, and Y is I or ClO₄.

173. The pyrylium compound according to Claim 165, represented by the following formula [10]:




[10],

wherein X is O, S, Se, or Te, and Y- is an anion.

174. The pyrylium compound according to Claim 173, wherein X is O or S, and Y is I or ClO₄.

175. The pyrylium compound according to Claim 165,

11




[11],

wherein X is O, S, Se, or Te, and Y- is an anion.

176. The pyrylium compound according to Claim 175, wherein X is O or S, and Y is I or ClO₄.

177. The pyrylium compound according to Claim 165,
represented by the following formula [12]:




[12],

wherein X is O, S, Se, or Te, and Y⁻ is an anion.

178. The pyrylium compound according to Claim 177, wherein X is O or S, and Y is I or ClO₄.


179. The pyrylium compound according to Claim 165, represented by the following formula [13]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

180. The pyrylium compound according to Claim 179, wherein X is O or S, and Y is I or ClO₄.


181. The pyrylium compound according to Claim 165, represented by the following formula [14]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

182. The pyrylium compound according to Claim 181, wherein X is O or S, and Y is I or ClO₄.

183. The pyrylium compound according to Claim 165, represented by the following formula [15]:



wherein X is O, S, Se, or Te, and Y⁻ is an anion.

184. The pyrylium compound according to Claim 183, wherein X is O or S, and Y is I or ClO₄.

185. The pyrylium compound according to Claim 165, wherein at least one hydrophilic group is introduced into at least one substituent of said pyrylium compound.

0994015.034704

ADD
b1